Imaging as a Biomarker: Standards for Change Measurements in Therapy 14-15 September 2006, NIST, Gaithersburg, MD

The Osteoarthritis Initiative (OAI): A Public-Private Partnership

Charles Peterfy, M.D., Ph.D.
Chief Medical Officer
Synarc, Inc.
San Francisco, CA

Osteoarthritis is a Major Unmet Medical Need

- Highly prevalent, chronic, debilitating and costly
 - 8% of the US population (21 M people) have physician-diagnosed OA
 - Many more have undiagnosed or subclinical OA
 - 50% >65 years: frequent knee pain and X-ray OA
 - But, only 1 in 6 consult a physician
 - By 2020, 20% (70 M people) >65 y
 - ⇒ 35 M Americans with X-ray OA
 - >200,000 total knee replacement procedures for OA
- Only symptomatic therapy
- No preventative or disease-modifying therapies (DMOADs) with proven efficacy and safety



Barriers to OA Drug Development

- Inadequate understanding of OA pathophysiology
 - Link between structural abnormalities and clinical outcomes
 - Difficulty identifying potentially modifiable therapeutic targets
- Epidemiological studies and clinical trials to elucidate these have to be large, long, difficult and expensive
- Existing cohorts offer valuable info, but are too small and inaccessible
- Lack of biomarkers which are suitably valid, discriminative and feasible

The Osteoarthritis Initiative: A Public-Private Partnership



Academia

NIH funded contracts
Access to public database and specimens

Industry

Study planning
Scientific input
Financial backing
Data & specimens

<u>Government</u>

NIH: Financial input
Contract funding
Holder of public database
Specimen repository

<u>FDA</u>



Goals of OAI

- Produce a public research resource
 - Identify and validate imaging, biochemical and genetic biomarkers for OA
 - Characterize early natural history and progression of OA
 - Evaluate risk and prognostic factors (including biomarkers) for OA onset and progression
- Main focus on knee OA
 - Limited data on hip and hand OA



Approach to Meet OAI Goals

- Prospective cohort study of knee OA
 - Diverse and representative sample of people with knee
 OA or at risk of developing knee OA
 - Serial measurements (annual)
 - Imaging of knee, hip and hand
 - Biospecimens for biochemical and genetic markers
 - Clinical assessment (joint Sx, disability) and risk factors
 - Public data archive on the web
 - Archived images for distribution to users
 - Archived biospecimens available by application



Government Partners

- National Institute of Arthritis & Musculoskeletal and Skin Diseases (NIAMS)
 - Gayle Lester (Project Officer)
- National Institute on Aging (NIA)
- National Center for Complementary & Alternative Medicine
- National Center on Minority Health & Health Disparities
- National Institute of Biomedial Imaging and Bioengineering
- National Institute of Dental & Craniofacial Research
- Office of Research on Women's Health
- Office of the Director, NIH



- Pharmaceutical Partners
 - GlaxoSmithKline
 - Christine Dabrowski and others
 - Merck & Company, Inc.
 - Chan Beals and others
 - Novartis Pharmaceuticals Corporation
 - Monica Luchi, and others
 - Pfizer, Inc
 - Marie Pierre Hellio LeGraverand-Gastineau, and others
 - Provide funding and participate in scientific decisions

- Clinical Centers
 - University of Maryland / Johns Hopkins University
 - Marc Hochberg, Joan Bathon
 - Memorial Hospital Rhode Island / Brown University
 - Charles Eaton, Timothy McAlindon
 - Ohio State University
 - Rebecca Jackson
 - University of Pittsburgh
 - Kent Kwoh
 - Subject recruitment and enrollment
 - 3.0 T Siemens Trio MR scanner installed at each center, dedicated to OAI

- Data Coordinating Center
 - University California San Francisco (Michael Nevitt)
- Imaging QA Center
 - Synarc, Inc. (Charles Peterfy)
- Steering Committee
- Observational Study Monitoring Board (OSMB)
- Consultants and Subcontractors



Imaging Work Group

- F. Boada
- W. Carter
- B. Dardzinski
- F. Eckstein
- J. Evelhoch
- D. Felson
- G. Gold
- B. Goodpaster
- M-P. Hellio Le Graverand
- M. Kothari
- P. Lang

- G. Lester
- M. Nevitt
- C. Peterfy
- T. Pellas
- C. Resnick
- D. Robertson
- E. Schneider
- L. Sharma
- G. Tung
- T. Woodworth

Progression and Incidence Cohorts

Number

1,330

Progression:

Sx knee OA at baseline



Follow for progression (Sx, function, structure)

3,215

Incidence:

Risk factors for Sx knee OA



Follow for incident Sx knee OA and other endpoints (X-ray OA)

200

"Nonexposed" control group:
No Sx, no X-ray OA,
No risk factors

190 incidents expected




Subjects: All Cohorts

- Inclusion
 - Men and Women
 - Ages 45 -79 years
 - All ethnic minorities (19%)
- Exclusion
 - Contraindications to MRI
 - History of inflammatory arthritis
 - RA Sx and medication screen
 - Bilateral end-stage disease (severe JSN and/or bilat TKR)



Examination Schedule



* 50% Progression cohort get 18 and 30 mo visits (MRI, biospecimens, Sx)



Designing the Imaging Protocol


- Provide imaging data on as many joint structures and features believed to be relevant to OA as possible
- Provide images able to support as broad a range of existing and anticipated measurement methods for each structure and feature as possible
- Balance scientific requirements for image quality and consistency against practical needs of high throughput and patient retention



1. Which articular structures and OA features to focus?

- Articular Cartilage
- Osteophytes
- Bone marrow edema
- Menisci
- Synovial effusion
- Cruciate and collateral ligaments
- Subarticular bone cysts
- Periarticular bursa
- Thigh muscles

OMERACT-OARSI Workshop for Consensus on OA Imaging, Bethesda, Dec 2002



- What measurement methods (quantitative and semi-quantitative) to support for each?
 - Most established existing biomarkers:
 - Cartilage volume / thickness (regional)
 - Semiquantitative scoring of cartilage and other structures
 - e.g., WORMS
 - Should also support as many future analyses as possible



- 3. Subject tolerance
 - 60 min 75 min
- 4. Generalizability of the results
 - Availability of hardware and pulse sequences
 - Commercial systems and pulse sequences
 - Standardized vs. diverse systems and pulse sequences
 - Identical, dedicated, state of the art systems at each site
 - Standardized protocol
 - ⇒3T MRI (Siemens Trio): ↑ resolution / speed, avoid obsolescence
 - Extrapolation of 3T data to 1.5T data?
 - Could not borrow protocols established for 1.5T



5. Future utility

Images must support broadest range of analyses possible now and tomorrow



Image Quality Control

Goals

- Uniform high-quality artifact-free images from all sites
- Longitudinal consistency of key parameters
 - SNR, CNR, signal homogeneity, spatial distortion
- Comparability of images from all 4 sites
- Minimum need for repeat imaging by preemptive correction of slowly developing problems



Image Quality Control

- Imaging procedure manuals
- Central and local training of imaging personnel
 - Technologists
 - Local radiologists
- On-going technical feedback
- Site visits to qualify imaging facilities




Image Quality Control

- Acceptance testing
- MRI system performance monitoring
 - Phantoms (ACR and OAI)
 - Regular performance tests


- Image quality
 - Check for protocol adherence
 - Critical image quality criteria
 - Anatomical coverage
 - Artifacts
 - Feedback to sites







Public Data Release http://w.w.w.oai.ucsf.edu/datarelease/



Public Data Release http://w.w.w.oai.ucsf.edu/datarelease/

- General information about OAI and its design
- Description of study data, procedures and materials
- On-line access to forms, operating manuals and data documentation
- Limited on-line data exploration
- Clinical data sets available for download to registered users
- Images distributed on various electronic media
- Biospecimens available by application to NIAMS-administered Biospecimen Resource Allocation Committee




Image Analyses

- OAI will make available a limited number of image assessments to address broad questions
 - Relationship between baseline structural and biochemical markers and risk factors with:
 - Onset or progression of X-ray JSW and MRI cartilage markers
 - Onset or progression of symptoms and disability
 - Relationship between changes in structural and biochemical markers with:
 - Onset or progression of X-ray JSW and MRI cartilage markers
 - Onset or progression of symptoms and disability



Image Analyses

- Baseline radiographic assessments that were done by readers at the clinical sites to assign subcohorts
 - Presence of definite tibiofemoral osteophytes (OARSI 1-3)
 - Mild-moderate (OARSI 1-2) vs. servere (OARSI 3) medial tibiofemoral joint-space narrowing



Image Analyses

- Central assessments of Progression subcohort TBD by Steering Committee and contracted out
 - Knee X-ray:
 JSW and subjective score at BL and FU -->



Knee MRI:
 cartilage volume / thickness at BL and FU
 semiquantitative score of OA features at BL



- Hand X-ray:
 presence and severity of OA at BL
- Pelvis X-ray: presence and severity of OA at BL



The Osteoarthritis Initiative

- OAI represents a new research model:
 - Creation of public resources:
 - database, images & specimen repository
- Scientific partnership:
 - Academic and industry scientists
- Novel funding mechanism for research:
 - Involvement of multiple companies and NIH to pool resources
- Other consortia have since been established

